

Current Situation on the Excess Mortality of COVID-19 for Various Countries

Bando H^{a,b}

^aTokushima University / Medical Research, Tokushima, Japan

^bIntegrative Medicine Japan (IMJ), Shikoku Island division, Tokushima, Japan

Article Info

Article History:

Received: 22 May, 2021

Accepted: 24 May, 2021

Published: 27 May, 2021

***Corresponding author:** Bando H,
Tokushima University, Medical
Research, Tokushima, Japan; Tel: +81-
90-3187-2485; E-mail:

pianomed@bronze.ocn.ne.jp

DOI:

<https://doi.org/10.36266/GJIDIT/112>

Abstract

For the research of COVID-19, the excess mortality has been adequate indicator for comparison of various situation from international point of view. The data of excess mortality in many countries were reported in Jan and April 2021. In current results for annual statistic, excess mortalities are US 458,000, Italy 89,100, England 85,400, Spain 84,100, Poland 60,100, and New Zealand 2500. Previous data showed Taiwan -4800, Australia -4700, and Japan -15000. For COVID-19 pandemic, previous data of excess deaths from public report shows the possibility of underestimation. Several factors are presumably involved in COVID-19 impact including public health measures and influenza prevalence.

Keywords: COVID-19; Excess Mortality; Health and Human Services (HHS); New Zealand; Japan

Copyright: © 2021 Bando H. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Commentary

Concerning COVID-19, several projects were found for predicting death rate, such as calculation of age-related relative risk (RR) [1] and fatality rates [2]. The excess mortality has been adequate indicator for the research of COVID-19. The international data from 77 countries were previously summarized and reported, in which some representative data as US 420,000, Brazil 170,000, England 80,000, Italy 64,000, Germany 30,000, New Zealand -2,100, Taiwan -4,800 and Japan -15,000 [3]. The author reported detail situation of Japan with minus excess mortality from several points of view as well as future perspectives for human resources and life [4]. The estimation of excess mortality is based on the additive hazard model, which separates observed mortality hazard into expected mortality and excess period. For example, the result from analysis in UK showed that absolute excess risk was 703 per 100,000 person-years (95% CI 687-719) for about one third of a year. In 2018, excess risk was 902 for one year. Thus, the risk of 1/3 year in 2020 equals to that of 3/4 year [5]. Most recent data of excess mortality were summarized for 29 high income countries [6]. Highest countries are US, Italy, UK, Spain and Poland, while New Zealand showed lower result. Overall men showed much higher than women. Both results of Jan and April 2020 are compared (Table 1) [3,4,6]. In this article, several current reports are introduced.

For COVID-19 pandemic, previous data of excess deaths from public report seemed to be underestimated. Then detail analysis was conducted by using the Joinpoint regression program [7]. Deaths in US was 1,336,561 during Mar-Aug 1, 2020, which was 20% more than expected. Out of 225,530 excess deaths,

150,541 cases (67%) seemed to be attributed to COVID-19. In recent report, excess deaths were 2,801,439 during Mar-Dec 2000, which showed 22.9% increase than expected with 522,368 excess deaths [8]. Among them, three subgroups were analyzed, where non-Hispanic Black, non-Hispanic White, Hispanic population showed 208.4, 157.0, 139.8 deaths per 100,000, respectively. Further, the excess death ratio was 16.9%, 61.1%, 16.7%, respectively. COVID-19 has brought remarkable increase in all-cause deaths in the US, which are mostly elder adults. In contrast, adults from 25-44 years were investigated [9]. The methods applied the data from Health and Human Services (HHS) in US. The results showed that all-cause deaths were 76,088 for Mar-July 2020 that was 11,899 more than expected 64,189 deaths (incident rate ratio 1.19). Covid-19 deaths were 4,535, which was 38% of measured excess mortality. For HHS region including New York and New Jersey showed incident rate was 2.30. Age-specific mortality rate was compared between US and European countries. As a result, adverse mortality situations in the US showed 400,700 excess in 2017 [10].

Several studies are from European countries. Excess mortality for care home resident in UK was evaluated from Jan 2017 to Aug 2020 [11]. The results showed that all care home was 6.5%, nursing home was 8.4% and residential home was 4.6% out of 29,542 excess deaths. Among them, 64.7% of excess deaths were suspected or confirmed by COVID-19. Odds ratio for experiencing COVID-19 was 1.8 for nursing services, 5.5 for older people and/or with dementia. From Italy, relationship between care home facilities and excess deaths was conducted [12]. As a result, significant higher excess death was found for the presence of care homes in the municipality. Especially, this situation seems to be driven from elderly people of 70 years and older. The influence of

COVID-19 was studied for 31 nursing homes (NH) in Italy [13]. Totally 3,946 NH residents showed females 73.9% and 87 years in median were included. Among them 1,136 was positive for COVID-19. As a result, the risk of death was increased as HR 1.85 by COVID-19.

Table 1: Excess mortality metrics for the countries in the dataset.

| Country | Estimated number of excess deaths in 2020 | |
|-----------------|---|-----------------|
| | Islam ref.6 | Karlinsky ref.3 |
| | April 29,2021 | Jan 27,2021 |
| United States | 458000 | 420000 |
| Italy | 89100 | 64000 |
| England & Wales | 85400 | 80000 |
| Spain | 84100 | 78000 |
| Poland | 60100 | 67000 |
| France | 43500 | 52000 |
| Germany | 25900 | 30000 |
| Belgium | 17900 | 18000 |
| Hungary | 16600 | 11000 |
| Netherlands | 15300 | 18000 |
| Czecho Republic | 14400 | |
| Sweden | 9300 | 9400 |
| Portugal | 8500 | 12000 |
| Greece | 7700 | 4700 |
| Austria | 6800 | 8600 |
| Scotland | 6800 | |
| Switzerland | 6800 | 6400 |
| Lithuania | 6800 | 6400 |
| Slovakia | 4400 | 2500 |
| South Korea | 4000 | -2500 |
| Slovenia | 3200 | 2800 |
| Nothern Ireland | 2200 | |
| Finland | 1000 | 890 |
| Latvia | 820 | 1700 |
| Estonia | 670 | 620 |
| Norway | -70 | -20 |
| Denmark | -160 | 650 |
| New Zealand | -2500 | -2100 |
| Mexico | | 270000 |
| Russia | | 270000 |
| Brazil | | 170000 |
| South Africa | | 96000 |
| Peru | | 93000 |
| Australia | | -4700 |
| Taiwan | | -4800 |
| Japan | | -15000 |

Israel is known to conduct early vaccination. Excess all-cause mortality rates were studied for Mar-Nov 2020 compared with 2017-2019 [14]. Total mortality rate was significantly 6% higher, where 11%, 13%, 19% higher in Aug, Sept, and Oct, 2020. Excess mortality was significantly 7% higher in 65-74/75-84 years, and 8% higher in 85≤ years, and greater for males than females in all ages. Ecological analysis was performed for 22 countries compared with 5-year historical all-cause mortality. There were excess deaths of 716,616 compared to last 5 years, where 64.3% of them were

attributed to COVID-19. Consequently, deaths data would be underestimated by at least 35% by COVID-19 [15]. Excess mortality data show minus number in New Zealand, Taiwan and Japan. New Zealand conducted strict intervention of public health during middle March for 12th week of 2020, such as early border closure, compulsory self-isolation after travelling, nationwide lockdown, limited contacts and isolation of infected cases [16]. In combination with the geographical isolation, these effective interventions brought COVID-19 elimination in early June, 2020 for 24th week [17]. Furthermore, persistent reduced mortality would be also from the absence of influenza epidemic in 2020. Consequently, both of public health measures and influenza prevalence were presumably involved in COVID-19 elimination [18]. In Taiwan, COVID-19 affected confirmed 911 cases, and 8 death until Jan 31, 2021 [19]. For statistic method, spline-smoothed Poisson regression models were used for death rate and seasonal effect from Jan 2015- Dec 2019. Then, expected deaths were calculated for Jan 2020 to Jan 2021, and excess deaths were estimated for -2,433 (95% CI: -10,715 to 5,850) [20]. This method also enables the analysis of various cause of reduction in mortality from pneumonia and influenza [21].

For Japan, healthcare access during COVID-19 impact was investigated for actual medical practice [22]. As a result, limited accessibility was found and future proposal for urgent supportive management against prolonged pandemics will be proposed. In summary, currently meaningful reports were described and discussed. The information will become hopefully useful for further research development. Conflict of interest: The author declares no conflict of interest.

References

- Banerjee A, Pasea L, Harris S, Gonzalez-Izquierdo A, Torralbo, A Shallcross L, et al. Estimating excess 1-year mortality associated with the COVID-19 pandemic according to underlying conditions and age: a population-based cohort study. *Lancet*. 2020; 395: 1715-1725.
- Verity R, Okell LC, Dorigatti I, Winskill P, Whittaker C, Imai N, et al. Estimates of the severity of coronavirus disease 2019: a model-based analysis. *Lancet Infect Dis*. 2020; 20: 669-677.
- Karlinsky A, Kobak D. The World Mortality Dataset: Tracking excess mortality across countries during the COVID-19 pandemic. *Med Rxiv*. 2021; 29: 1-10.
- Bando H. Fewer Annual Deaths in Japan with Negative Excess Mortality for COVID-19. *Asp Biomed Clin Case Rep*. 2021; 4: 84-87.
- Joy M, Hobbs FDR, McGagh D, Akinyemi O, Lusignan S. Excess mortality from COVID-19 in an English sentinel network population. *Lancet Infect Dis*. 2021; 21: 74.
- Islam N, Shkolnikov VM, Acosta RJ, Klimkin I, Kawachi I, Irizarry RA, et al. Excess deaths associated with covid-19 pandemic in 2020: age and sex disaggregated time series analysis in 29 high income countries. *BMJ* 2021; 1137-1137.
- Woolf SH, Chapman DA, Sabo RT, Weinberger DM, Hill L, Taylor DDH. Excess Deaths from COVID-19 and Other Causes, March-July 2020. *JAMA*. 2020; 324: 1562-1564.
- Woolf SH, Chapman DA, Sabo RT, Zimmerman EB. Excess Deaths from COVID-19 and Other Causes in the US, March 1, 2020, to January 2, 2021. *JAMA*. 2021; 325: 1786-1789.
- Faust JS, Krumholz HM, Du C, Mayes KD, Lin Z, Gilman C, et al. All-Cause Excess Mortality and COVID-19-Related Mortality Among US Adults Aged 25-44 Years, March-July 2020. *JAMA*. 2021; 325: 785-787.
- Preston SH, Vierboom YC. Excess mortality in the United States in the 21st century. *Proc Natl Acad Sci U S A*. 2021; 118: 2024850118.
- Morciano M, Stokes J, Kontopantelis E, Turner JT. Excess mortality for care home residents during the first 23 weeks of the COVID-19 pandemic in England: a national cohort study. *BMC Med*. 2021; 19: 71.
- Alacevich C, Cavalli N, Giuntella O, Lagravinese R, Moscone F, Nicodemo C. The presence of care homes and excess deaths during the COVID-19 pandemic: Evidence from Italy. *Health Econ*. 2021.
- Veronese N, Koyanagi A, Stangherlin V, Mantoan P, Chiavalin M, Tudor F, et al. Mortality attributable to COVID-19 in nursing home residents: a retrospective study. *Aging Clin Exp Res*. 2021; 24: 1-7.
- Haklai Z, Aburbeh M, Goldberger N, Gordon ES. Excess mortality during the COVID-19 pandemic in Israel, March-November 2020: when, where, and for whom?. *Isr J Health Policy Res*. 2021; 10: 17.
- Kung S, Doppen M, Black M. Underestimation of COVID-19 mortality during the pandemic. *ERJ Open Res*. 2021; 7: 00766-2020.
- Baker MG, Wilson N, Anglemeyer A. Successful Elimination of Covid-19 Transmission in New Zealand. *N Engl J Med*. 2020; 383: 56.
- Kung S, Doppen M, Black M, Hills T, Kearns N. Reduced mortality in New Zealand during the COVID-19 pandemic. *Lancet*. 2021; 397: 25.
- Hills T, Kearns N, Kearns C, Beasley R. Influenza control during the COVID-19 pandemic. *Lancet*. 2020; 396: 1633-1634.
- Taiwan Centers for Disease Control Press Releases on January, 2021.
- Chen EY, Sachs MC, Dickman PW. No evidence of substantial underreporting of COVID-19 deaths in Taiwan during 2020. *J Formos Med Assoc*. 2021; 24.
- Gao W, Sanna M, Huang G, Hefler M, Tsai MK, Wen CP. Examining Population Health During the COVID-19 Pandemic: All-Cause, Pneumonia and Influenza, and Road Traffic Deaths in Taiwan. *Ann Intern Med*. 2021.
- Makiyama K, Kawashima T, Nomura S. Trends in Healthcare Access in Japan during the First Wave of the COVID-19 Pandemic, up to June 2020. *Int J Environ Res Public Health*. 2021; 18:3 271.